

Acknowledgments.—I thank Youli Fu, a middle school teacher in Changpo Town, Danzhou City; Shiping Gong, my graduate student in the Academy of Life Science, Beijing Normal University; Riheng Song, my undergraduate in the Department of Biology, Hainan Normal University; and my friend Hua Li; we all took part in this activity together. I especially thank James Ford Parham for help, including helping me revise my English. Finally, I thank the National Natural Science Foundation of China (No. 30260019), Foundation for University Key Teacher by the Ministry of Education, the Oregon Zoo Foundation, the Chicago Zoo Foundation, Ross Kiester, and Theodore Papenfuss for supporting conservation and research on turtles in Hainan.

LITERATURE CITED

- BUSKIRK, J.R. 1993. Captive propagation and husbandry of the Vietnamese leaf turtle (*Geoemyda spengleri*). *Vivarium* 5(3):28-33.
- CHAU, L., FELLOWES, J., AND LAM, V. (Eds.). 2002. Records of globally and nationally threatened wildlife in Guangdong, Guangxi and Hainan. *Living Forests - Newsletter for Biodiversity Conservation in South China* 4:30-32.
- DE BRUIN, R.W.F. AND ARTNER, H.G. 1999. On the turtles of Hainan Island, southern China. *Chelonian Conservation and Biology* 3(3):479-486.
- ERNST, C.H. AND BARBOUR, R.W. 1989. *Turtles of the World*. Washington, DC: Smithsonian Institution Press, 313 pp.
- GRESSITT, J.L. 1940. On a collection of amphibians and reptiles made on Hainan Island. *Peking Natural History Bulletin* 15:175-193.
- IVERSON, J.B. 1992. *A Revised Checklist with Distribution Maps of the Turtles of the World*. Richmond, IN: Privately printed, 363 pp.
- LI, Z.Y. 1958. Report on the investigation of reptiles of Hainan Island. *Chinese Journal of Zoology* 2(4):234-239.
- LAU, M. AND SHI, H. 2000. Conservation and trade of terrestrial and freshwater turtles and tortoises in the People's Republic of China. In: van Dijk, P.P., Stuart, B.L., and Rhodin, A.G.J. (Eds.). *Asian Turtle Trade: Proceedings of a Workshop on Conservation and Trade of Freshwater Turtles and Tortoises in Asia*. *Chelonian Research Monographs* 2:30-38.
- PETZOLD, H.-G. 1984. *Cuora galbinifrons* and other Southeast Asian turtles in the Berlin Zoological Gardens. *CDC Newsl.* 3(1-4):4-7.
- RHODIN, A.G.J. 2002. Conservation and trade of freshwater turtles and tortoises in Asia: review of status and threats using IUCN Red List and CITES criteria. *CITES Technical Workshop on Trade in Freshwater Turtles and Tortoises in Asia*. Kunming, China, March 2002, Doc. 3.2, 45 pp.
- SCHMIDT, K.P. 1927. The reptiles of Hainan. *Bulletin of the American Museum of Natural History* 54:395-465.
- SHI, H. 2000. A brief introduction to "International workshop on the conservation and trade of freshwater turtles and tortoises in Asia". *Sichuan Journal of Zoology* 19(3):189.
- SICHUAN INSTITUTE OF BIOLOGY (ZHAO, E. AND YANG, S.). 1976. An introduction to Order Chelonia. *Materials for Herpetological Research*, Chengdu 3:61-68.
- TIAN, W. AND JIANG, Y. (Eds.). 1986. *Identification Manual of Chinese Amphibians and Reptiles*. Beijing: Science Press, 142 pp.
- YAO, W. AND LIU, Z. 1995. Preliminary observation on *Geoemyda spengleri*, a record new to Anhui Province. *Chinese Journal of Zoology* 30(3):22-24.
- ZHANG, M., ZONG, Y., AND MA, J. 1998. *Fauna Sinica. Reptilia Vol. 1. General Accounts of Reptilia. Testudoformes and Crocodiliformes*. Beijing: Science Press, 213 pp.
- ZHAO, E. 1997. Studies on taxonomy and distribution of Chinese chelonians. In: Zhao, E. (Ed.). *Chinese Chelonian Research. Chinese Society for the Study of Amphibians and Reptiles, Herpetological Series No. 9, Sichuan Journal of Zoology* 15 (Suppl.):1-26.
- ZHAO, E. 1998. *China Red Data Book of Endangered Animals. Vol. 3. Amphibia and Reptilia*. Beijing: Science Press, 330 pp.
- ZHAO, E. AND ADLER, K. 1993. *Herpetology of China. Society for the Study of Amphibians and Reptiles, Contributions to Herpetology, No. 10*, 522 pp.

Received: 15 January 2003

Revised and Accepted: 1 November 2004

Chelonian Conservation and Biology, 2005, 4(4):956-959
© 2005 by Chelonian Research Foundation

✓ Occurrence of Diamondback Terrapins, *Malaclemys terrapin*, on Bermuda: Native or Introduced?

JOHN DAVENPORT^{1,2}, ANNE F. GLASSPOOL³,
AND LISA KITSON^{1,3}

¹Department of Zoology, Ecology and Plant Science, and
²Environmental Research Institute, University College Cork,
Lee Maltings, Prospect Row, Cork, Ireland

[Fax: 353-21-4-270562; E-mail: jdavenport@zoology.ucc.ie];

³Bermuda Aquarium, Natural History Museum and Zoo,
Flatts, FLBX, Bermuda

[E-mails: bamzcare@ibl.bm; biobabe@ibl.bm]

ABSTRACT.—A breeding population of the American diamondback terrapin, *Malaclemys terrapin*, occurs on the oceanic island of Bermuda; most of the population lives in two brackish lakes surrounded by mangrove vegetation. The question of whether the species reached Bermuda from the USA naturally or by deliberate or inadvertent introduction is considered.

The emydid diamondback terrapin, *Malaclemys terrapin*, is widely distributed in estuaries and salt marshes on the east coast of the USA from Texas to Massachusetts. During the past decade it has been discovered that there is a breeding population of *M. terrapin* on the isolated island of Bermuda (32°18'N, 64°46'W), some 960 km from the US coast (Fig. 1). The recent colonization of many bodies of water in Bermuda by released pet specimens of the American red-eared slider (*Trachemys scripta elegans*) has resulted in closer scrutiny of Bermudian pools, so greater knowledge of the distribution of diamondback terrapins on Bermuda is now available. This paper presents preliminary distributional information, and also considers the question of whether diamondback terrapins have reached Bermuda by natural means or by human intervention.

The currently accepted native Bermudian herpetofauna is limited to the endemic skink, *Eumeces longirostris* (the




Figure 1. Map to show position of Bermuda in relation to the USA. Large arrows indicate southern (Corpus Christi, Texas) and northern (Cape Cod, Massachusetts) distributional limits of *Malaclemys terrapin*.

only endemic terrestrial tetrapod), and four species of marine turtles. However, there have been numerous well-documented and dated introductions of amphibians and reptiles (several small lizard species of the genus *Anolis*, the cane toad, *Bufo marinus*, and two species of tree frog, as well as the red-eared slider). No such documentation exists for *M. terrapin*. Although present-day Bermuda is now an essentially suburban country (Sterrer, 1998) with a resident population of about 60,000 people and numerous tourists (0.5 million/yr), there is therefore the possibility that the island has harbored an extra, but unrecognized, native reptile for a considerable period of time.

Results. — Diamondback terrapins were first reported in the mid-1990s to the Bermuda Aquarium, Natural History Museum and Zoo (BAMZ) by staff of a local golf course, the Mid Ocean Club, who found eggs in sand bunkers. The eggs

were incubated at BAMZ and hatchlings released at the golf course in two small brackish man-made ponds that clearly contained numerous (between 10 to 100) mature male and female diamondback terrapins, as well as specimens of *T. s. elegans*.

Further investigation by BAMZ staff revealed that terrapins were also present in nearby Mangrove Lake, a relatively large natural brackish water body. They have been reported from all parts of the lake, which is completely surrounded by thick mangrove vegetation and separated from the sea by a narrow strip of land. However, the numbers present are currently unknown and an effective mark-recapture census would be a substantial undertaking and logistically difficult. Terrapins are also known from neighboring Trott Pond, another natural brackish pond. Local residents report that terrapins have been present in Mangrove Lake at least for several decades. It seems probable that the artificial golf course ponds were colonized by terrapins from one or other of the much larger natural ponds.

We recently collected three adult female diamondback terrapins from the pools at Mid Ocean Club (carapace lengths 197, 213, 215 mm) (Fig. 2). Their sizes were quite close to maximum recorded lengths for female diamondback terrapins (ca. 225 mm). All had extremely smooth carapaces with negligible scute grooves and rather indistinct markings. The dorsal keel was of negligible height. Carapace color was variable. One animal (215 mm) had a light brown shell, with the most recent scute growth being ivory in color. Its carapace was essentially elliptical and the marginal scutes showed only limited upcurling posteriorly. The skin was pale gray with dark gray spots. There was no moustache stripe. The other two had oval olive green shells, with recent scute growth being light green. Both had finely spotted light green to gray skin and no moustache. Their




Figure 2. Photographs of diamondback terrapins, *Malaclemys terrapin*, collected recently on Bermuda.

shells showed slightly more marginal upcurling than the brown animal.

Discussion. — Bermuda and its surrounding archipelago is one of the most remote island systems in the world. However, it is known that the Gulf Stream, which runs close to the North American coast between Florida and Cape Hatteras, spins off eddies that pass regularly through Bermudian waters. These eddies have been implicated in the transport of postlarval reef fish from Florida waters to Bermuda (Glasspool, 1994) and also in the transport (probably from Florida) of the extinct Bermudian tortoise *Hesperotestudo bermudae* to the islands, where it occurred some 300,000 years ago (Meylan and Sterrer, 2000). These authors estimated that the tortoise would have floated more than 2000 km to reach Bermuda, possibly achieving the passage in as little time as a week or two.

Among emydid turtles, diamondback terrapins are almost uniquely suited to long-distance travel at sea. Their estuarine natural habitats are often highly saline for weeks at a time (Bentley et al., 1967; Gilles-Baillien, 1970; Dunson, 1970, 1976, 1985). They are capable of living for long periods in sea water because they have low skin permeability to salts and water, excrete salts from orbital salt glands, accumulate urea in the blood plasma (thus minimizing osmotic water loss), and have behavioral responses that prevent drinking and minimize eating when in sea water for long periods (Schmidt Nielsen and Fänge, 1958; Bentley et al., 1967; Gilles-Baillien, 1970; Dunson, 1970, 1976, 1985; Davenport and Macedo, 1990; Davenport and Ward, 1993). Most freshwater emydids would succumb to osmotic dehydration and hyperchloremia within a few days in sea water.

Diamondback terrapins are well-suited to the conditions in which they live in Bermuda today. There are no surface streams on the island and the rock formations are highly permeable to sea water. In consequence, most pools are brackish rather than fresh, with the precise salinity depending to a large extent on recent rainfall. It is known that *Malaclemys* is capable of exploiting transient sources of rainwater (Davenport and Macedo, 1990) and is probably the best-adapted brackish-water turtle so far studied. There is little doubt that diamondback terrapins *could* have reached Bermuda naturally.

However, *did* diamondback terrapins reach Bermuda naturally? This is a much more difficult question to answer. *A priori*, diamondback terrapins could a) have been imported as pets and subsequently released, b) been imported as food animals, perhaps during the terrapin culture 'craze' of the early 20th century, or c) have arrived without human intervention as described above.

There is no evidence that diamondback terrapins have ever been imported as pets to Bermuda. Outside the USA they have never been a conspicuous part of the herpetofauna pet trade.

From the 1880s through to the 1920s in the USA, there was great demand for diamondback terrapins as gourmet

food items. Initially satisfied by capture of wild terrapins, by 1900 demand, plus declining stocks, had triggered the establishment of ranches and farms, particularly at Beaulieu, Georgia, and Beaufort, North Carolina (Gadow, 1901; Coker, 1906; Hildebrand and Hatsel, 1926; Hildebrand, 1932). Terrapins were not only eaten in the USA, but were exported to Europe and Brazil. The terrapin trade largely collapsed with the 1930s recession in the USA, though cook books published terrapin recipes well into the 1950s. There is no oral or written record of any attempt to farm terrapins in Bermuda, though it is conceivable that someone might have imported a few animals for consumption, but released them instead.

Given the scenario for possible natural colonization outlined above, it would seem likely that a possible natural source population for Bermudian terrapins would have to be in the southern part of the species' range, perhaps in Florida (the apparent source of the endemic skink *Eumeces longirostris* and the extinct fossil *Hesperotestudo bermudae* according to Meylan and Sterrer, 2000). There is no fossil evidence of long-term Bermudian residence for *Malaclemys*. A skeleton (with some scutes intact) was found in a dry cave near Harrington Sound (close to the two natural ponds in which terrapins are currently found). Carbon dating of such recent material would probably be of limited use unless it was medieval or older. Genetic analysis of Bermudian terrapins has not been carried out, and would in any case have to be related to the genetic makeup of diamondback terrapins from the populations living along the eastern coast of the USA. The species' wide geographical distribution, perhaps coupled with limited gene flow between populations living in brackish water habitats separated by open coast, has led to an unusual degree of diversity, with seven subspecies currently recognized (Carr, 1952; Pritchard, 1979).

If Bermudian *M. terrapin* proved to be genetically identical with *M. terrapin terrapin* (the northern diamondback) a natural origin for the terrapins would be virtually ruled out because their distribution is too far north to permit sea transport by the Gulf Stream to Bermuda. A more credible candidate, *M. t. centrata* (the Carolina diamondback) is difficult to separate from *M. t. terrapin* on morphological grounds as there are "all sorts of variations and intergradations" (Carr, 1952). To complicate matters further, farmed terrapins were a mixture of the two subspecies (Carr, 1952) and were released when farms closed. The features of the three terrapins we collected (Fig. 2) are not compatible with published descriptions (Pritchard, 1979) of *M. t. macrospilota*, *M. t. pileata*, or *M. t. littoralis*, all of which have deeply sculpted shells and high dorsal keels. The absence of a moustache and the presence of concentric shell markings exclude *M. t. tequesta*. Of the remaining three subspecies, the smooth, marbled carapace seems more compatible with descriptions of *M. t. centrata* than *M. t. terrapin*. However, it is difficult to rule out *M. t. rhizophorarum*, as that subspecies is relatively poorly studied, also has a smooth carapace with

slight dorsal keel, and occurs in extreme southern Florida where the Gulf Stream sweeps by close to shore.

Conclusions and Recommendations. — The Bermudian population of *M. terrapin* appears to be the only breeding population outside the USA. It is apparently thriving, though a full population estimate would be desirable to establish a benchmark for future study. Whether the species is native to Bermuda or introduced is of conservation and management significance. If native, then full legislative protection and appropriate conservation measures would be appropriate. If introduced, then study of the impact of terrapins upon the brackish-water fauna of Bermuda would be sensible.

The present investigation does not provide conclusive support for either hypothesis, and it is probable that a natural introduction could only be firmly proved if skeletal remains of pre-colonial age emerge at some time in the future. However, a natural introduction could be disproved if the Bermudian specimens can be shown to have closer affinities with *M. t. terrapin* than either *M. t. centrata* or *M. t. rhizophorarum*. There appears not to be an adequate genetic library presently available to permit this to be achieved by DNA analysis. Hopefully this situation will be clarified in the future.

Acknowledgments. — This is Contribution #73, Bermuda Biodiversity Project (BBP), Bermuda Aquarium, Natural History Museum and Zoo.

LITERATURE CITED

- BENTLEY, P.J., BRETZ, W.L., AND SCHMIDT-NIELSEN, K. 1967. Osmoregulation in the diamondback terrapin, *Malaclemys terrapin centrata*. *Journal of Experimental Biology* 46:161-167.
- CARR, A.F. 1952. *Handbook of Turtles. The Turtles of the United States, Canada, and Baja California*. Ithaca, NY: Cornell University Press, 542 pp.
- COKER, R.E. 1906. The cultivation of the diamond-back terrapin. *Bulletin of the North Carolina Geological Survey* 14:1-67.
- DAVENPORT, J. AND MACEDO, E.A. 1990. Behavioural osmotic control in the euryhaline diamondback terrapin *Malaclemys terrapin* Latreille: responses to low salinity and rainfall. *Journal of Zoology (London)* 220(3):487-496.
- DAVENPORT, J. AND WARD, J.F. 1993. The effects of salinity and temperature on appetite in the diamondback terrapin *Malaclemys terrapin* (Latreille). *Herpetological Journal* 3(3):95-98.
- DUNSON, W.A. 1970. Some aspects of electrolyte and water balance in three estuarine reptiles, the diamondback terrapin, American and "salt water" crocodiles. *Comparative Biochemistry and Physiology* 32:161-174.
- DUNSON, W.A. 1976. Salt glands in reptiles. In: Gans, C. and Dawson, W.R. (Eds.). *Biology of the Reptilia*. Vol. 5, Physiology A. London: Academic Press, pp. 413-445.
- DUNSON, W.A. 1985. Effect of water salinity and food salt content on growth and sodium efflux of hatchling diamondback terrapins (*Malaclemys*). *Physiological Zoology* 58(6):736-747.
- GADOW, H. 1901. *Amphibia and Reptiles*. London: Macmillan and Co.
- GILLES-BAILLIEN, M. 1970. Urea and osmoregulation in the diamondback terrapin *Malaclemys centrata centrata* (Latreille). *Journal of Experimental Biology* 52:691-697.
- GLASSPOOL, A.F. 1994. Larval distribution, population structure and gene flow in Bermuda's reef fish. Ph.D. Thesis, University of Wales, UK.
- HILDEBRAND, S.F. 1932. Growth of the diamond-back terrapins: size attained, sex ratio and longevity. *Zoologica* 9(15):551-563.
- HILDEBRAND, S.F. AND HATSEL, C. 1926. Diamond-back terrapin culture at Beaufort, North Carolina. U.S. Bureau of Fisheries Economic Circular No. 60, 20 pp.
- MEYLAN, P.A. AND STERRER, W. 2000. *Hesperotestudo* (Testudines: Testudinidae) from the Pleistocene of Bermuda, with comments on the phylogenetic position of the genus. *Zoological Journal of the Linnean Society* 128:51-76.
- PRITCHARD, P.C.H. 1979. *Encyclopedia of Turtles*. Neptune, NJ: TFH Publications, 895 pp.
- SCHMIDT-NIELSEN, K. AND FANGE, R. 1958. Salt glands in marine reptiles. *Nature* 182:783-785.
- STERRER, W. 1998. How many species are there in Bermuda? *Bulletin of Marine Science* 62(3):809-840.

Received: 9 April 2003

Revised and Accepted: 29 March 2004